

COHESIVENESS IN PROMISE PROBLEMS *

ULRIKE BRANDT AND HERMANN K.-G. WALTER¹

Abstract. Promise problems have been introduced in 1985 by S.Even e.a. as a generalization of decision problems. Using a very general approach we study solvability and unsolvability conditions for promise problems of set families and languages. We show, that cores of unsolvability are completely determined by partitions of cohesive sets. We prove the existence of cores in unsolvable promise problems assuming certain closure properties for the given set family. Connections to immune sets and complexity cores are presented. Furthermore, results about cohesiveness with respect to the language families from the Chomsky hierarchy are given.

1991 Mathematics Subject Classification. 68Q45.

INTRODUCTION

In 1985 S.Even, A.L.Selman and Y.Yacobi [4] introduced the concept of *promise problems* as a generalization of decision problems. A promise problem consists of a pair of disjoint sets A and B with $A, B \subseteq S$ and a given set family $\mathcal{F} \subseteq 2^S$, where S is some basic (usually infinite) set. (A, B) is *solvable for \mathcal{F}* if a $Q \subseteq S$ exists with $Q \in \mathcal{F}$ and $Q^c \in \mathcal{F}$ and $A \subseteq Q$ and $B \subseteq Q^c$, where Q^c is the complement of Q in S . In the case $B = A^c$ (A, B) is a *decision problem*. In applications $S = X^*$, where X is a finite nonempty alphabet and $\mathcal{F} = \mathcal{L}$ is language family or a complexity class $\mathcal{F} = \mathcal{C}$. From an algorithmic point of view considering a promise problem (A, B) an algorithm may only produce a Yes-answer for all instances $x \in A$ and a No-answer for all $x \in B$, while no decisive answer is expected for $x \notin A \cup B$. Solvability of promise problems can be linked to the existence of approximation or "special case" algorithms (see [2]). Thus with respect to complexity of algorithms a more refined look than for decision problems is possible. Promise problems have been considered for various fields of algorithmic computations. Especially, some decision problems which are difficult to solve allow efficient algorithm once they are weakened to a promise problem. The reader can find an overview in [5]. Looking at the theory

* We acknowledge seminal discussions with MARTIN ZIEGLER.

¹ Fachbereich Informatik, Technische Universität Darmstadt, Germany

of recursive functions [8], the separation principle is a precursor of the concept of promise problems. Furthermore, we can use the notion of cohesive sets, also known in the theory of recursive function, by an appropriate generalization. This turns out to be the characterizing indicator for (un-)solvability of promise problems. It was well-known, that only set-theoretic arguments can be used in dealing with these concepts. We mention especially the theorem of Dekker-Myhill [8] which asserts the existence of cohesive sets under very weak conditions. Our main results are a theorem about the existence of unsolvability cores for an unsolvable promise problem (A, B) and the characterization of unsolvability cores via cohesiveness of $A \cup B$. The latter enables us, to study the influence of closure operations on the unsolvability of promise problems. Though the existence of cohesive sets is guaranteed under very mild conditions, it is quite difficult, to exhibit cohesive languages with nice properties. We determine cohesive sets and noncohesive sets for language families from the Chomsky hierarchy and for families given by number theoretic properties. Especially, we prove a structure result for alphabets X with two or more letters. For some special cases we can at least assert the existence of recursive cohesive languages. Using results from [2], the connection to complexity cores gives a similar result for recursive language families and complexity classes. We assume the reader to be familiar with the theory of recursive functions and sets (see [2], [8], [9]) and standard theory of formal languages (see [6], [7]).

Our study of promise-problems was proposed to us by M. Ziegler who raised the question answered in theorem 5.13.

1. SET- AND LANGUAGE FAMILIES - BASIC NOTATIONS AND RESULTS

In the following a *basic* set S is given and we assume for set families $\mathcal{F} \subseteq \mathbf{2}^S$. Moreover, sets $A, A', B, B', C, \dots, Q, \dots$ are always subsets of S and *singletons* $\{s\}$ are identified with s . We mainly deal with denumerable set families \mathcal{F} ; i.e. a function $\mathbf{e}_{\mathcal{F}} : \mathbb{N}_0 \rightarrow \mathbf{2}^S$ with $\mathbf{e}_{\mathcal{F}}(\mathbb{N}_0) = \mathcal{F}$ exists (*enumeration* of \mathcal{F}). Consider the *boolean* operations *union*, *intersection* and *complementation* in connection with *set families* \mathcal{F} . The boolean operations can be lifted to binary operations between set families \mathcal{F}_1 and \mathcal{F}_2 and unary operations for \mathcal{F} . Define $\mathcal{F}_1 \oplus \mathcal{F}_2 = \{A \cup B | A \in \mathcal{F}_1 \text{ and } B \in \mathcal{F}_2\}$, $\mathcal{F}_1 \odot \mathcal{F}_2 = \{A \cap B | A \in \mathcal{F}_1 \text{ and } B \in \mathcal{F}_2\}$ and the closure operations

$$\begin{aligned} \mathcal{F}^{\mathbf{u}} &= \{A_1 \cup \dots \cup A_n | n \geq 1, A_i \in \mathcal{F} \text{ for } 1 \leq i \leq n\}(\text{union}), \\ \mathcal{F}^{\mathbf{s}} &= \{A_1 \cap \dots \cap A_n | n \geq 1, A_i \in \mathcal{F} \text{ for } 1 \leq i \leq n\}(\text{intersection}), \\ \mathcal{F}^{\mathbf{co}} &= \{A^c | A \in \mathcal{F}\}, \mathcal{F}^{\mathbf{cc}} = \mathcal{F} \cup \mathcal{F}^{\mathbf{co}}, (\text{complementation}) \text{ and} \\ \mathcal{F}^{\mathbf{b}} &= ((\mathcal{F}^{\mathbf{cc}})^{\mathbf{s}})^{\mathbf{u}}(\text{boolean closure}). \end{aligned}$$

Moreover, we will frequently use $\mathcal{F}^{\mathbf{dc}} = \mathcal{F} \cap \mathcal{F}^{\mathbf{co}}$.

Note, that $(\mathcal{F}^{\mathbf{u}})^{\mathbf{s}} = (\mathcal{F}^{\mathbf{s}})^{\mathbf{u}}$ (*distributivity*), $(\mathcal{F}^{\mathbf{co}})^{\mathbf{u}} = (\mathcal{F}^{\mathbf{s}})^{\mathbf{co}}$ (*deMorgan*), $(\mathcal{F}^{\mathbf{cc}})^{\mathbf{dc}} = \mathcal{F}^{\mathbf{cc}}$ and $(\mathcal{F}^{\mathbf{co}})^{\mathbf{co}} = \mathcal{F}$. There are numerous (mostly trivial) relations between these operations, for example.

Proposition 1.1. Let $\mathcal{F}_1, \mathcal{F}_2 \subseteq \mathcal{2}^S$.

- (1) $\mathcal{F}_1 \oplus \mathcal{F}_2 \subseteq \mathcal{F}_1 \Rightarrow \mathcal{F}_1 \oplus \mathcal{F}_2^u \subseteq \mathcal{F}_1$ and $\mathcal{F}_1^u \oplus \mathcal{F}_2 \subseteq \mathcal{F}_1^u$.
- (2) $\mathcal{F}_1 \odot \mathcal{F}_2^{co} \subseteq \mathcal{F}_1 \Rightarrow \mathcal{F}_1 \odot (\mathcal{F}_2^{co})^u \subseteq \mathcal{F}_1^u$ and $\mathcal{F}_1^u \odot \mathcal{F}_2^{co} \subseteq \mathcal{F}_1^u$.

In the following we frequently use the combined operation of *variation of \mathcal{F}* by \mathcal{V} defined by $\mathcal{F} \pm \mathcal{V} = \mathcal{F} \oplus \mathcal{V} \cup \mathcal{F} \odot \mathcal{V}^{co}$.

Proposition 1.2. Let $\mathcal{F}, \mathcal{V} \subseteq \mathcal{2}^S$ with $\mathcal{V} \neq \emptyset$ and $\mathcal{F} \pm \mathcal{V} \subseteq \mathcal{F}$.

- (1) $\emptyset, S \in \mathcal{F} \Leftrightarrow \mathcal{V}^{cc} \subseteq \mathcal{F}$.
- (2) $\mathcal{F}^{cc} \pm \mathcal{V} \subseteq \mathcal{F}^{cc}, \mathcal{F}^u \pm \mathcal{V} \subseteq \mathcal{F}^u, \mathcal{F}^s \pm \mathcal{V} \subseteq \mathcal{F}^s$ and $\mathcal{F}^b \pm \mathcal{V} \subseteq \mathcal{F}^b$.
- (3) $\mathcal{F} \pm \mathcal{V}^u \subseteq \mathcal{F}$

In the case $\mathcal{V} = \mathbf{fin}(S) = \{A \subseteq S \mid A \text{ finite}\}$, the condition $\mathcal{F} \pm \mathbf{fin}(S) \subseteq \mathcal{F}$ is just the closure under *finite variation*. Note that, $\mathbf{fin}(S)^{cc} = (\mathbf{fin}(S)^{cc})^b$ and $\mathcal{F} \odot \mathbf{fin}(S) \subseteq \mathbf{fin}(S)$. By prop.1.2.(2) $\mathcal{F}^{cc}, \mathcal{F}^u, \mathcal{F}^s, \mathcal{F}^b$ are closed under finite variation, if \mathcal{F} is closed under finite variation.

Consider the case $S = X^*$, where X is a nonempty, finite alphabet and X^* is the *free monoid over X* . As usual $L \subseteq X^*$ is called a *language* and $\mathcal{L} \subseteq \mathcal{2}^{X^*}$ a *language family*. The elements of X^* are the *words* $w = x_1 \dots x_n$ ($x_i \in X$ for $1 \leq i \leq n$) and the *empty word* $\mathbf{1}$. The *length* of w is $|w| = n$ and $|\mathbf{1}| = 0$. *Concatenation* " wv " of words is the monoid operation with identity $\mathbf{1}$. The operation can be lifted to $\mathcal{2}^{X^*}$. For $L_{1,2}$ the *complex product* is defined by $L_1 L_2 = \{w_1 w_2 \mid w_1 \in L_1, w_2 \in L_2\}$. L^* is the *generated submonoid*.

On X^* we can define various (partial) orderings. The following two ones are of interest to us. Define for $v, w \in X^*$ the *prefix-ordering* by $w \leq v(\mathbf{pref}) \Leftrightarrow v \in wX^*$. Given a bijection $\mathbf{ord} : X \rightarrow [0 \dots b-1]$ ($b = \#(X)$) we can define also a well-ordering \mathbf{lex}_{ord} by $w \leq v(\mathbf{lex}_{ord})$ if and only if $|w| < |v|$ or $\forall u \in X^*, x, y \in X : ux \leq w(\mathbf{pref})$ and $uy \leq v(\mathbf{pref}) \Rightarrow \mathbf{ord}(x) \leq \mathbf{ord}(y)$. Since \mathbf{lex}_{ord} is a well-ordering, we can define a successor function \mathbf{succ}_{ord} for $w \in X^*$ by $\mathbf{succ}_{ord}(w) = \mathbf{min}\{v \in X^* \mid w \neq v \text{ and } w \leq v(\mathbf{lex}_{ord})\}$, where the minimum is taken with respect to \mathbf{lex}_{ord} . Then $\mathbf{char}^*(i) = \mathbf{succ}_{ord}^i(\mathbf{1})$ ($i \geq 0$) defines a bijection $\mathbf{char}^* : \mathbb{N}_0 \rightarrow X^*$.

The language families from the Chomsky hierarchy are $\mathcal{L}_{r.e.}(X)$ (*recursively enumerable languages*), $\mathcal{L}_{cs}(X)$ (*contextsensitive languages*), $\mathcal{L}_{cf}(X)$ (*contextfree languages*) and $\mathcal{L}_{reg}(X)$ (*regular languages*). All these families are closed under variation by $\mathcal{L}_{reg}(X)$. By encoding the *generating grammars* we find special enumerations $\mathbf{e}_{r.e.}, \mathbf{e}_{cs}, \mathbf{e}_{cf}$ and \mathbf{e}_{reg} of the corresponding language family. With these enumerations we can study decision problems and constructions for the *descriptive devices* (grammars). Look for example at the *word-problem* for $\mathcal{L}_{cs}(X)$. Using $0, 1 \in \mathbb{N}_0$ as truth values, define the predicate $\mathbf{word}_{cs}(i, j) = \text{"char}^*(i) \in \mathbf{e}_{cs}(j)\text{"}$ ($i, j \geq 0$). Then $\mathbf{word}_{cs} \in \mathbf{rec}_2$, where \mathbf{rec}_n ($n \geq 0$) is the set of *n-ary recursive functions*. In the case of *complexity classes* \mathcal{C} we can find enumerations $\mathbf{e}_{\mathcal{C}}$, such that $\mathbf{word}_{\mathcal{C}}(i, j) = \text{"char}^*(i) \in \mathbf{e}_{\mathcal{C}}(j)\text{"}$ ($i, j \geq 0$) is recursive. Here we have to use as descriptive devices Turingmachines with reasonable resource bounds (*time-/space-constructibility* (see [2])). More general, let

$\mathbf{word}_e(i, j) = \text{"char}^*(i) \in \mathbf{e}(j)\text{"}$ ($i, j \geq 0$) for any $\mathbf{e} : \mathbb{N}_0 \rightarrow \mathbf{2}^{X^*}$. We call \mathbf{e} *WP-recursive* if and only if $\mathbf{word}_e \in \mathbf{rec}_2$. A language family \mathcal{L} is *WP-recursive*, if a WP-recursive enumeration \mathbf{e} of \mathcal{L} exists. In this case $\mathcal{L} \subseteq \mathcal{L}_{\mathbf{rec}}(X) = \mathcal{L}_{\mathbf{r.e.}}(X)^{\mathbf{dc}}$ (*recursive languages*). Note, that for WP-recursive families a *uniform* solution for the word-problem exists. Complexity classes are WP-recursive and closed under variation by $\mathcal{L}_{\mathbf{reg}}(X)$.

Considering $\mathcal{L}_{\mathbf{reg}}(X)$ and $\mathcal{L}_{\mathbf{cf}}(X)$ we obtain by the classical decidability results, that the predicates $\mathbf{empty}_{\mathbf{cf}}(i) = \text{"e}_{\mathbf{cf}}(i) = \emptyset\text{"}$ and $\mathbf{finite}_{\mathbf{cf}}(i) = \text{"e}_{\mathbf{cf}} \in \mathbf{fin}(X^*)\text{"}$ ($i \geq 0$) are recursive. Moreover, $f_{\mathbf{sect}} \in \mathbf{rec}_2$ (*intersection with regular sets*) and $f_{\mathbf{comp}} \in \mathbf{rec}_1$ (*complementation of regular sets*) exist with $\mathbf{e}_{\mathbf{cf}}(i) \cap \mathbf{e}_{\mathbf{reg}}(j) = \mathbf{e}_{\mathbf{cf}}(f_{\mathbf{sect}}(i, j))$ and $\mathbf{e}_{\mathbf{reg}}(i)^c = \mathbf{e}_{\mathbf{reg}}(f_{\mathbf{comp}}(i))(i, j \geq 0)$. Using all these functions, we find $\mathbf{incl}(i, j) = \text{"e}_{\mathbf{cf}}(i) \subseteq \mathbf{e}_{\mathbf{reg}}(j)\text{"} = \mathbf{empty}_{\mathbf{cf}}(f_{\mathbf{sect}}(i, f_{\mathbf{comp}}(j)))(i, j \geq 0)$, hence $\mathbf{incl} \in \mathbf{rec}_2$.

In the following, at various points we are faced with marking languages at the left, i.e. we have to consider the *left translation* " wL ".

Proposition 1.3. *For all languages $L, L_{1,2}$ and $w \in X^*$:*

- (1) $w(L_1 \cup L_2) = wL_1 \cup wL_2$ and $w(L_1 \cap L_2) = wL_1 \cap wL_2$,
- (2) $wL^c = (wL)^c \cap wX^*$ and $(wL)^c = wL^c \cup (wX^*)^c$.

For a language family \mathcal{L} define $\mathcal{L}^{\mathbf{ltr}} = \{wL \mid w \in X^*, L \in \mathcal{L}\}$ (*left translation*). $\mathcal{L}^{\mathbf{ltr}}$ is another closure operation and $\mathcal{L} = \mathcal{L}^{\mathbf{ltr}}$ if and only if $xL \in \mathcal{L}$ for any $x \in X$. Moreover, a companion to proposition 1.2.(1) (with $\mathcal{V} = \mathbf{fin}(X^*)$) holds.

Proposition 1.4. *Let $\mathcal{L} = \mathcal{L}^{\mathbf{u}} = \mathcal{L}^{\mathbf{ltr}}$. Then $\mathbf{fin}(X^*)^{\mathbf{cc}} \subseteq \mathcal{L}$ if and only if $\emptyset, \mathbf{1}, X^* \in \mathcal{L}$.*

Proof. Let $\emptyset, \mathbf{1}, X^* \in \mathcal{L}$. Since $w = w\mathbf{1}$ and $\mathcal{L} = \mathcal{L}^{\mathbf{ltr}}$, singletons are in \mathcal{L} . But then $\mathbf{fin}(X^*) \subseteq \mathcal{L}$, because $\mathcal{L} = \mathcal{L}^{\mathbf{u}}$. Let $X^k = \{w \in X^* \mid |w| = k\}$ ($k \geq 0$). Then X^k is finite. Hence $X^k X^*$ is the finite union of sets wX^* , i.e. $X^k X^* \in \mathcal{L}^{\mathbf{u}} = \mathcal{L}$. Let $L \in \mathbf{fin}(X^*)$ and $k > \mathbf{max}\{|w| \mid w \in L\}$, then $L^c = ((X^k X^*)^c \setminus L) \cup X^k X^*$. Note that $((X^k X^*)^c \setminus L)$ is finite and therefore an element of \mathcal{L} as shown before. In total $L^c \in \mathbf{fin}(X^*) \oplus \mathcal{L} \subseteq \mathcal{L}^{\mathbf{u}} = \mathcal{L}$, i.e. $\mathbf{fin}(X^*)^{\mathbf{co}} \subseteq \mathcal{L}$. \square

In connection with boolean operations we get

Lemma 1.5. *If $\mathcal{L} = \mathcal{L}^{\mathbf{ltr}}$, then*

- (1) $(\mathcal{L}^{\mathbf{u}})^{\mathbf{ltr}} = \mathcal{L}^{\mathbf{u}}$ and $(\mathcal{L}^{\mathbf{s}})^{\mathbf{ltr}} = \mathcal{L}^{\mathbf{s}}$.
- (2) $(\mathcal{L}^{\mathbf{cc}})^{\mathbf{ltr}} = \mathcal{L}^{\mathbf{cc}}$, if additionally $\mathcal{L} \pm \mathcal{L}_{\mathbf{reg}}(X) \subseteq \mathcal{L}$.

Proof. (1) By prop.1.3.(1): $(\mathcal{L}^{\mathbf{u}})^{\mathbf{ltr}} \subseteq (\mathcal{L}^{\mathbf{ltr}})^{\mathbf{u}} = \mathcal{L}^{\mathbf{u}}$. Analogously, $(\mathcal{L}^{\mathbf{s}})^{\mathbf{ltr}} \subseteq \mathcal{L}^{\mathbf{s}}$. (2) Since $wX^* \in \mathcal{L}_{\mathbf{reg}}(X)$ for all $w \in X^*$, we get by our assumption, prop.1.3.(2) and prop.1.2.(2) $(\mathcal{L}^{\mathbf{co}})^{\mathbf{ltr}} \subseteq (\mathcal{L}^{\mathbf{ltr}})^{\mathbf{co}} \odot \mathcal{L}_{\mathbf{reg}}(X) \subseteq (\mathcal{L}^{\mathbf{ltr}})^{\mathbf{cc}} \pm \mathcal{L}_{\mathbf{reg}}(X) = \mathcal{L}^{\mathbf{cc}} \pm \mathcal{L}_{\mathbf{reg}}(X) \subseteq \mathcal{L}^{\mathbf{cc}}$. This shows $(\mathcal{L}^{\mathbf{cc}})^{\mathbf{ltr}} \subseteq \mathcal{L}^{\mathbf{cc}}$. \square

Next we look at the inverse of left translations (removing left markers).

Definition 1.6. \mathcal{L} is *ltr-cancellative* if and only if for all $L \subseteq X^*$ and $w \in X^*$: $wL \in \mathcal{L} \Rightarrow L \in \mathcal{L}$.

Proposition 1.7. *If \mathcal{L} is ltr-cancellative, then*

- (1) \mathcal{L}^u is ltr-cancellative
- (2) \mathcal{L}^{co} is ltr-cancellative, if additionally $\mathcal{L} \pm \mathcal{L}_{\text{reg}}(X) \subseteq \mathcal{L}$.

Proof. (1) Let $wL = L_1 \cup \dots \cup L_n$ with $L_i \in \mathcal{L}$ for $1 \leq i \leq n$. Then each $L_i \subseteq wX^*$, i.e. $L_i = wL'_i$. Since \mathcal{L} is ltr-cancellative, $L'_i \in \mathcal{L}$. But then $L = L'_1 \cup \dots \cup L'_n \in \mathcal{L}^u$.
(2) If $wL \in \mathcal{L}^{\text{co}}$, then $(wL)^c \in \mathcal{L}$. Since $(wL)^c = wL^c \cup (wX^*)^c$ by prop.1.3.(2) and $\mathcal{L} \pm \mathcal{L}_{\text{reg}}(X) \subseteq \mathcal{L}$, we get $wL^c \in \mathcal{L}$ and therefore $L^c \in \mathcal{L}$. Hence $L \in \mathcal{L}^{\text{co}}$, i.e. \mathcal{L}^{co} is ltr-cancellative. \square

All families from the Chomsky hierarchy and all complexity classes are ltr-cancellative and closed under left translation.

2. COHESIVENESS

Definition 2.1. A is \mathcal{F} -cohesive ($A \in \mathbf{cohesive}(\mathcal{F})$) if and only if $A \notin \mathbf{fin}(S)$ and for any $B \in \mathcal{F}^{\text{dc}}$: $(A \cap B \notin \mathbf{fin}(S) \Rightarrow A \cap B^c \in \mathbf{fin}(S))$.

Remark: The definition of cohesiveness given in §12.3 of [8] is equivalent to $\mathcal{L}_{\text{r.e.}}(X)^{\text{cc}}$ -cohesiveness.

Directly from the definition we get

Proposition 2.2.

- (1) $\mathbf{cohesive}(\mathcal{F}) = \mathbf{cohesive}(\mathcal{F}^{\text{co}}) = \mathbf{cohesive}(\mathcal{F}^{\text{dc}})$
- (2) $\mathcal{F}_1 \subseteq \mathcal{F}_2 \Rightarrow \mathbf{cohesive}(\mathcal{F}_2) \subseteq \mathbf{cohesive}(\mathcal{F}_1)$.

Next we study the influence of closure operations on cohesiveness. Especially, we discuss boolean operations and closure under finite variation and left translation.

Lemma 2.3. *If $\mathcal{F} = \mathcal{F}^{\text{cc}}$, then $\mathbf{cohesive}(\mathcal{F}) = \mathbf{cohesive}(\mathcal{F}^{\text{b}})$.*

Proof. By our assumption $\mathcal{F}^{\text{b}} = (\mathcal{F}^{\text{s}})^u$. Consider $A \in \mathbf{cohesive}(\mathcal{F})$ and $B \in \mathcal{F}^{\text{b}}$ with $A \cap B \notin \mathbf{fin}(S)$. Then $B = B_1 \cup \dots \cup B_n$ with $B_i \in \mathcal{F}^{\text{s}}$ for $1 \leq i \leq n$. Now, $A \cap (B_1 \cup \dots \cup B_n) = (A \cap B_1) \cup \dots \cup (A \cap B_n) \notin \mathbf{fin}(S)$. But then $C = B_j$ exists with $A \cap C \notin \mathbf{fin}(S)$. Since $C \subseteq B$, we know that $A \cap B^c \subseteq A \cap C^c$. Hence, if $A \cap C^c \in \mathbf{fin}(S)$, then $A \cap B^c \in \mathbf{fin}(S)$.

Since $C \in \mathcal{F}^{\text{s}}$, $C = C_1 \cap \dots \cap C_m$ with $C_i \in \mathcal{F}$ for $1 \leq i \leq m$. Furthermore, $A \cap C \notin \mathbf{fin}(S)$, so that for every i , $A \cap C_i \notin \mathbf{fin}(S)$. But then, by the cohesiveness of A , $A \cap C_i^c \in \mathbf{fin}(S)$ for $1 \leq i \leq m$ and therefore $A \cap C^c = A \cap (C_1 \cap \dots \cap C_m)^c = A \cap (C_1^c \cup \dots \cup C_m^c) = (A \cap C_1^c) \cup \dots \cup (A \cap C_m^c) \in \mathbf{fin}(S)$. \square

Proposition 2.4. $\mathbf{cohesive}(\mathcal{F})$ is closed under finite variation.

Proof. Consider $A \in \mathbf{cohesive}(\mathcal{F})$, $C \in \mathbf{fin}(S)$ and some $B \in \mathcal{F}$. Assume that $(A \cup C) \cap B = (A \cap B) \cup (C \cap B) \notin \mathbf{fin}(S)$. Since $C \cap B \in \mathbf{fin}(S)$, $A \cap B \notin \mathbf{fin}(S)$ and therefore $A \cap B^c \in \mathbf{fin}(S)$ due to the cohesiveness of A . Since $C \cap B^c \in \mathbf{fin}(S)$ as well, $(A \cup C) \cap B^c = (A \cap B^c) \cup (C \cap B^c) \in \mathbf{fin}(S)$.

In the second step, assume that $(A \cap C^c) \cap B = (A \cap B) \cap C^c \notin \mathbf{fin}(S)$. Then $A \cap B \notin \mathbf{fin}(S)$, i.e. $A \cap B^c \in \mathbf{fin}(S)$, because A is \mathcal{F} -cohesive. But then $(A \cap C^c) \cap B^c = (A \cap B^c) \cap C^c \in \mathbf{fin}(S)$. \square

For $S = X^*$ and left translation we can show

Lemma 2.5. *If \mathcal{L} is ltr-cancellative, $\mathcal{L} = \mathcal{L}^{\text{ltr}}$ and $\mathcal{L} \pm \mathcal{L}_{\text{reg}}(X) \subseteq \mathcal{L}$, then for any $L \in \mathcal{L}, w \in X^* : L \in \mathbf{cohesive}(\mathcal{L})$ if and only if $wL \in \mathbf{cohesive}(\mathcal{L})$.*

Proof. Suppose $L \in \mathbf{cohesive}(\mathcal{L})$, i.e. $L \cap B \notin \mathbf{fin}(S)$ implies $L \cap B^c \in \mathbf{fin}(S)$ for any $B \in \mathcal{L}$. Consider $w \in X^*$ and $B \in \mathcal{L}^{\text{dc}}$ with $wL \cap B \notin \mathbf{fin}(S)$. Then we have to show, that $wL \cap B^c \in \mathbf{fin}(S)$. Clearly, $wL \cap B = w(L \cap A)$ for A with $wA = B \cap wX^*$. Thus, $A \in \mathcal{L}$, since \mathcal{L} is ltr-cancellative and closed under finite variation by regular sets. By the same arguments we get $A^c \in \mathcal{L}$, too: To see this, observe that $wA^c = (wA)^c \cap (wX^*)$ by prop.1.3.(2) and therefore $wA^c = (B \cap wX^*)^c \cap wX^* = (B^c \cup (wX^*)^c) \cap wX^* = B^c \cap wX^*$.

Now, $wL \cap B \notin \mathbf{fin}(S)$ implies $L \cap A \notin \mathbf{fin}(S)$. Hence, $L \cap A^c \in \mathbf{fin}(S)$ by the cohesiveness of \mathcal{L} . Since by prop.1.3.(1), $w(L \cap A^c) = wL \cap wA^c = wL \cap (B^c \cap wX^*) = wL \cap B^c$, $wL \cap B^c$ must be finite, too.

Conversely, suppose $wL \in \mathbf{cohesive}(\mathcal{L})$ for some $w \in X^*$. Fix $B \in \mathcal{L}^{\text{dc}}$ with $L \cap B \notin \mathbf{fin}(S)$. We show, that $L \cap B^c$ is finite. Observe, that $wB \in \mathcal{L}^{\text{dc}}$ by the closure properties of \mathcal{L} . Now, $L \cap B \notin \mathbf{fin}(X^*)$ implies $wL \cap wB^c \notin \mathbf{fin}(X^*)$ and therefore $wL \cap (wB)^c \in \mathbf{fin}(X^*)$, because wL is \mathcal{L} -cohesive. But $wL \cap (wB)^c = wL \cap wB^c = w(L \cap B^c)$, i.e. $L \cap B^c \in \mathbf{fin}(X^*)$. \square

The existence of cohesive sets for denumerable set families is guaranteed by a result of J. C. E. Dekker and J. Myhill (cf. Theorem VI in §12.3 of [8]).

Theorem 2.6. *(Dekker and Myhill) Let \mathcal{F} be an denumerable set family. Then for any $A \notin \mathbf{fin}(S)$ there is a subset B of A with $B \in \mathbf{cohesive}(\mathcal{F})$.*

The following fact is obvious:

Proposition 2.7. *If $B \subseteq A, B \notin \mathbf{fin}(S)$ and $A \in \mathbf{cohesive}(\mathcal{F})$ then $B \in \mathbf{cohesive}(\mathcal{F})$.*

A natural generalization of Theorem VII(ii) in §12.3 of [8] is

Lemma 2.8. *If $A, B \in \mathbf{cohesive}(\mathcal{F})$ and $A \cap B \notin \mathbf{fin}(S)$ then $A \cup B \in \mathbf{cohesive}(\mathcal{F})$*

Remark: Note that the condition " $A \cap B \notin \mathbf{fin}(S)$ " in le.2.8. is necessary. To see this, consider $X = \{a, b\}$ and \mathcal{L} satisfying the condition of le.2.5. If $L \in \mathbf{cohesive}(\mathcal{L})$, then $aL, bL \in \mathbf{cohesive}(\mathcal{L})$. But $(aL \cup bL) \cap aX^* = aL \notin \mathbf{fin}(X^*)$ and $bL \subseteq (aL \cup bL) \cap (aX^*)^c \notin \mathbf{fin}(X^*)$. Hence, $aL \cup bL \notin \mathbf{cohesive}(\mathcal{L})$.

Cohesiveness is a stronger condition than immunity for sets in connection with set families. For a set family \mathcal{F} a set A is defined to be \mathcal{F} -immune if it is infinite and has no infinite subset in \mathcal{F} , i.e. if $A^c \cap B \neq \emptyset$ for any $B \in \mathcal{F} \setminus \mathbf{fin}(S)$ (cf. e.g. [2], [8]). Let $\mathbf{immune}(\mathcal{F})$ denote the family of all \mathcal{F} -immune sets. Clearly, infinite subsets of \mathcal{F} -immune sets are \mathcal{F} -immune and $\mathbf{immune}(\mathcal{F}_2) \subseteq \mathbf{immune}(\mathcal{F}_1)$, if $\mathcal{F}_1 \subseteq \mathcal{F}_2$.

Proposition 2.9. *If \mathcal{F} is closed under finite variation and $A \in \mathbf{cohesive}(\mathcal{F}) \setminus \mathcal{F}$ then $A \in \mathbf{immune}(\mathcal{F})$.*

Proof. Suppose $B \in \mathcal{F} \setminus \mathbf{fin}(S)$ exists with $A^c \cap B = \emptyset$. Then $A \cap B = B \notin \mathbf{fin}(S)$ and therefore $A \cap B^c \in \mathbf{fin}(S)$, because A is \mathcal{F} -cohesive. Since \mathcal{F} is closed under finite variation $A = B \cup (A \cap B^c) \in \mathcal{F}$ - a contradiction. \square

Remark: Theorem V in §12.3 of [8] shows, that any $L \in \mathbf{cohesive}(\mathcal{L}_{\text{r.e.}}(X)^{\text{cc}})$ is not only immune but hyperimmune and even hyperhyperimmune.

Example 2.10. Let $X = \{a, b\}$. Then $A = \{a^n b^n | n > 0\}$ is $\mathcal{L}_{\text{reg}}(X)$ -immune (use the pumping lemma for $\mathcal{L}_{\text{reg}}(X)$), but not $\mathcal{L}_{\text{reg}}(X)$ -cohesive (consider e.g. $R = (a^2)^*(b^2)^*$).

Example 2.11. \mathcal{F} -cohesive languages need not necessarily be outside of \mathcal{L} :

- (1) Any $A \notin \mathbf{fin}(S)$ is $\mathbf{fin}(S)^{\text{cc}}$ -cohesive, for example $X^* \in \mathbf{fin}(X^*)^{\text{cc}}$.
- (2) By a theorem of Friedberg $L \in \mathcal{L}_{\text{r.e.}}(X)^{\text{co}}$ exists with $L \in \mathbf{cohesive}(\mathcal{L}_{\text{r.e.}}(X)^{\text{cc}})$ (see Theorem XI in 12.4 of [8] for details).

3. COHESIVENESS OF LANGUAGES

We derive special results for cohesiveness with respect to language families, especially for the families from the Chomsky hierarchy and complexity classes. For all these families \mathcal{L} any \mathcal{L} -cohesive language has a specific structural property. This property is connected to infinite words. Infinite words can be defined using **pref**-isotone and length-preserving functions.

Definition 3.1. $f : \mathbb{N}_0 \rightarrow X^*$ is *sequential* if and only if for any $n \geq 0$: $|f(n)| = n$ and $f(n) \leq f(n+1)$ (**pref**).

Lemma 3.2. If $\#(X) > 1$ and $L \in \mathbf{cohesive}(\mathcal{L}_{\text{reg}}(X))$, then a sequential $f_L : \mathbb{N}_0 \rightarrow X^*$ exists with $L \setminus f_L(n)X^* \in \mathbf{fin}(X^*)$ for any $n \geq 0$.

Proof. The key to the proof is the following

Assertion: If $L \in \mathbf{cohesive}(\mathcal{L}_{\text{reg}}(X))$, then for all $u, v \in X^*$ with $|u| = |v|$: $L \cap uX^*, L \cap vX^* \notin \mathbf{fin}(X^*)$ implies $u = v$.

Suppose the contrary, i.e. $u, v \in X^*$ exist with $|u| = |v|, L \cap uX^*, L \cap vX^* \notin \mathbf{fin}(X^*)$ and $u \neq v$. Then $uX^* \cap vX^* = \emptyset$. Hence, $vX^* \cap L \subseteq (uX^*)^c \cap L$ and therefore $(uX^*)^c \cap L \notin \mathbf{fin}(X^*)$. Thus, $L \notin \mathbf{cohesive}(\mathcal{L}_{\text{reg}}(X))$ - a contradiction.

Since $L \notin \mathbf{fin}(X^*)$, we can find to any $n \geq 0$ some $w \in X^*$ with $|w| = n$ and $L \cap wX^* \notin \mathbf{fin}(X^*)$. Define $f_L(n) = w$. By the assertion f_L is uniquely determined. Furthermore, if $u \leq w$ (**pref**), then $L \cap wX^* \subseteq L \cap uX^*$. Hence, $L \cap uX^* \notin \mathbf{fin}(X^*)$ and by the assertion $f_L(|u|) = u$. That is, f_L is sequential. Moreover, since $L \in \mathbf{cohesive}(\mathcal{L}_{\text{reg}}(X))$, $L \cap (f_L(n)X^*)^c \in \mathbf{fin}(X^*)$ for all $n \geq 0$. □

From the assertion in this proof we get additionally the following

Corollary 3.3. If $\#(X) > 1$ and $L \in \mathbf{cohesive}(\mathcal{L}_{\text{reg}}(X))$, then for any $L' \subseteq L$ with $L' \notin \mathbf{fin}(X^*)$: $f_L = f_{L'}$.

Proof. Suppose $n \geq 0$ exists with $f_L(n) \neq f_{L'}(n)$. We know by le.3.2. and prop.2.7. that $L' \cap f_{L'}(n)X^*, L \cap f_L(n)X^* \notin \mathbf{fin}(X^*)$. Furthermore, $L' \cap f_{L'}(n)X^* \subseteq L \cap f_{L'}(n)X^*$. Hence, $L \cap f_{L'}(n)X^*$ is infinite. This is a contradiction to the above assertion. □

Next, we focus our attention to $\mathcal{L}_{\text{reg}}(X)$ -cohesiveness in connection with $\mathcal{L}_{\text{cf}}(X)$ and $\mathcal{L}_{\text{cs}}(X)$. To $L \subseteq X^*$ we associate the *length-language* $|L| = \{a^{|w|} \mid w \in L\} = \lambda_x(L)$, where $\lambda_x(w) = |w| (w \in X^*)$. Define $\mathcal{L}_{\text{lreg}}(X) = \{L \subseteq X^* \mid |L| \in \mathcal{L}_{\text{reg}}(a)\}$. Note that $\{a^n b^n a^n \mid n \geq 0\} \in \mathcal{L}_{\text{lreg}}(\{a, b\})$.

Lemma 3.4. *If $L \in \mathcal{L}_{\text{lreg}}(X)$, then $L \notin \text{cohesive}(\mathcal{L}_{\text{reg}}(X))$.*

Proof. Since $L \notin \text{fin}(X^*)$, $|L| \in \mathcal{L}_{\text{reg}}(a) \notin \text{fin}(X^*)$. By the pumping lemma for $\mathcal{L}_{\text{reg}}(a)$ $\alpha > 0$ and $\beta \geq 0$ exist with $(a^\alpha)^* a^\beta \subseteq |L|$. Consider $R = (a^{2\alpha})^* a^\beta \in \mathcal{L}_{\text{reg}}(a) \setminus \text{fin}(a^*)$. Then $R \subseteq |L|$ and $R^c \cap |L| \notin \text{fin}(a^*)$. But $\lambda_x^{-1}(R), \lambda_x^{-1}(R^c) \in \mathcal{L}_{\text{reg}}(X)$, $L \cap \lambda_x^{-1}(R), L \cap \lambda_x^{-1}(R^c) \notin \text{fin}(X^*)$ and $\lambda_x^{-1}(R^c) \subseteq \lambda_x^{-1}(R)^c$. \square

Since $\mathcal{L}_{\text{reg}}(X) \subseteq \mathcal{L}_{\text{cf}}(X) \subseteq \mathcal{L}_{\text{lreg}}(X)$, we get immediately

Lemma 3.5. *If $L \in \text{cohesive}(\mathcal{L}_{\text{cf}}(X))$, then $L \notin \mathcal{L}_{\text{cf}}(X)$.*

Inspecting the "construction" from the proof of the Dekker-Myhill-theorem (Theorem VI in §12.3 of [8]) yields

Theorem 3.6. *For any $L \in \mathcal{L}_{\text{cf}}(X) \setminus \text{fin}(X^*)$ a recursive language $L' \subseteq L$ exists with $L' \in \text{cohesive}(\mathcal{L}_{\text{reg}}(X))$.*

Proof. We refine the proof of the Myhill-Dekker-theorem. Define inductively

$$\begin{aligned} L_0 &= L, \\ L_{n+1} &= \mathbf{if} \ L_n \cap \mathbf{e}_{\text{reg}}(n) \notin \text{fin}(X^*) \ \mathbf{then} \ L_n \cap \mathbf{e}_{\text{reg}}(n) \ \mathbf{else} \ L_n \cap \mathbf{e}_{\text{reg}}(n)^c \ \mathbf{fi} \\ &\quad (n \geq 0). \end{aligned}$$

Then for $n \geq 0$ $L_{n+1} \subseteq L_n$ and $L_n \in \mathcal{L}_{\text{cf}}(X)$. Moreover, $L_n \notin \text{fin}(X^*)$ and $L_n \notin \text{cohesive}(\mathcal{L}_{\text{reg}}(X))$ by le.3.5. Now, a function g exists with $L_{g(n+1)} \subseteq L_{g(n)} \subseteq L_n$. Define the function h by $h(n) = \min\{m \mid \text{char}^*(m) \in L_{g(n)} \setminus L_{g(n+1)}\}$ ($n \geq 0$). Let $L' = \text{char}^*(h(\mathbb{N}_0))$. The above mentioned proof of Theorem VI in §12.3 of [8] asserts $L' \in \text{cohesive}(\mathcal{L}_{\text{reg}}(X))$.

It remains to prove, that any step in this construction is computable. Let $\mathbf{e}_{\text{cf}}(i_0) = L$. Define

$$\begin{aligned} f(0) &= i_0, \\ f(n+1) &= \mathbf{if} \ \mathbf{finite}_{\text{cf}}(f_{\text{sect}}(f(n), n)) = 0 \ \mathbf{then} \ f_{\text{sect}}(f(n), n) \\ &\quad \mathbf{else} \ f_{\text{sect}}(f(n), f_{\text{comp}}(n)) \ \mathbf{fi} \ (n \geq 0). \end{aligned}$$

Since $\mathbf{finite}_{\text{cf}}$ is recursive, $f \in \text{rec}_1$ and $\mathbf{e}_{\text{cf}}(f(n)) = L_n (n \geq 0)$. Next, consider the predicate d defined by $d(n, m) = "(m > n) \text{ and } (L_n = L_m)"$. Then $d(n, m) = "(m > n) \text{ and } (\forall 1 \leq i \leq m - n : L_{n+i} = L_m)" = "(m > n) \text{ and } (\forall 0 \leq i \leq m - n : (L_{n+i} \subseteq \mathbf{e}_{\text{reg}}(n+i)) \text{ or } (L_{n+i} \subseteq \mathbf{e}_{\text{reg}}(n+i)^c))"$. Using \mathbf{incl} and f_{comp} we get $d \in \text{rec}_2$. By this the function $g'(n) = \min\{m \mid (m > n) \text{ and } d(f(n), m) = 0\}$ is recursive, as well. Now, with the help of f, g' the function g from above can be defined by $g(0) = f(0), g(n) = f(g'(n-1)) (n > 0)$, i.e. $g \in \text{rec}_1$, since $f, g' \in \text{rec}_1$. Since $w(m, n) = "\text{char}^*(m) \in L_{g(n)} \setminus L_{g(n+1)}" = \text{word}_{\text{cs}}(m, f(g(n)))$ is a

recursive function, $h(n) = \min\{m \mid \mathbf{char}^*(m) \in L_{g(n)} \setminus L_{g(n+1)}\} \in \mathbf{rec}_1$ and we can conclude that $L'' = \mathbf{char}^*(h(\mathbb{N}_0)) \in \mathcal{L}_{\mathbf{r.e.}}(X)$. But then an infinite $L' \subseteq L''$ exists with $L' \in \mathcal{L}_{\mathbf{rec}}(X)$. Since $L'' \in \mathbf{cohesive}(\mathcal{L}_{\mathbf{reg}}(X))$, $L' \in \mathbf{cohesive}(\mathcal{L}_{\mathbf{reg}}(X))$, too. \square

Dealing with $\mathcal{L}_{\mathbf{cs}}(X)$ we can use number-theoretic considerations. In the case $X = \{a\}$ we know $\mathcal{L}_{\mathbf{reg}}(X) = \mathcal{L}_{\mathbf{cf}}(X)$ and can use the pumping lemma for regular sets.

Lemma 3.7. (*number-problems*) *Let $X = \{a\}$.*

- (1) $L_{exp} = \{a^{2^n} \mid n > 0\} \notin \mathbf{cohesive}(\mathcal{L}_{\mathbf{reg}}(X))$ and $L_{exp} \in \mathcal{L}_{\mathbf{cs}}(X)$.
- (2) $L_{fac} = \{a^{n!} \mid n > 0\} \in \mathbf{cohesive}(\mathcal{L}_{\mathbf{reg}}(X))$ and $L_{fac} \in \mathcal{L}_{\mathbf{cs}}(X)$.

Proof. (1) Clearly, $2^{2k} \bmod 3 = (3+1)^k \bmod 3 = 1$ Hence $2^{2k+1} \bmod 3 = 2(2^{2k} \bmod 3) \bmod 3 = 2$. By this $\{a^{2^{2k}} \mid k \geq 0\} \subseteq a(a^3)^* = R$, while $\{a^{2^{2k+1}} \mid k \geq 0\} \subseteq R^c$. In total $L_{exp} \notin \mathbf{cohesive}(\mathcal{L}_{\mathbf{reg}}(X))$, because $L_{exp} \cap R, L_{exp} \cap R^c \notin \mathbf{fin}(a^*)$.

(2) Consider $R \in \mathcal{L}_{\mathbf{reg}}(X)$ with $L_{fac} \cap R \notin \mathbf{fin}(a^*)$. Using the pumping lemma for $\mathcal{L}_{\mathbf{reg}}(X)$ $\alpha > 0$ and β exist with $L_{fac} \cap a^\beta(a^\alpha)^* \notin \mathbf{fin}(a^*)$ and $a^\beta(a^\alpha)^* \subseteq R$. Due to the infinity of the intersection we can find some $y \geq \max(\alpha, \beta)$, such that $\alpha x + \beta = y!$ for some $x \in \mathbb{N}_0$. Since $y \geq \alpha$, α divides $y!$. Hence, β is an integer multiple of α , too and we get $\alpha x + \beta = \alpha(x + \beta') = y!$ for some $\beta' \leq y$. But then for any $y' \geq y$ a z exists with $\alpha(z + \beta') = y'!$, namely $z = (x + \beta') \frac{y'!}{y!} - \beta'$, and we get in total $L_{fac} \cap a^\beta(a^\alpha)^* = L_{fac} \setminus C$ for some finite set C and therefore $L_{fac} \cap R^c \in \mathbf{fin}(a^*)$. \square

A result similar to le.3.5. is

Proposition 3.8. *If $L \in \mathbf{cohesive}(\mathcal{L}_{\mathbf{rec}}(X))$, then $L \notin \mathcal{L}_{\mathbf{r.e.}}(X)$.*

Proof. Consider $L \in \mathbf{cohesive}(\mathcal{L}_{\mathbf{rec}}(X))$ and suppose that $L \in \mathcal{L}_{\mathbf{r.e.}}(X)$. Since L is infinite, $L' \in \mathcal{L}_{\mathbf{rec}}(X) \setminus \mathbf{fin}(X^*)$ exists with $L' \subseteq L$. Clearly, $L = L_1 \cup L_2$ with $L_{1,2} \in \mathcal{L}_{\mathbf{rec}}(X) \setminus \mathbf{fin}(X^*)$ and $L_1 \cap L_2 = \emptyset$. But then $L \cap L_1 = L_1 \notin \mathbf{fin}(X^*)$ and $L \cap L_2 \subseteq L \cap L_1^c \notin \mathbf{fin}(X^*)$ - a contradiction. \square

4. SOLVABILITY OF PROMISE PROBLEMS

Remember that given a set familie $\mathcal{F}(A, B)$ is a promise problem, if $A \cap B = \emptyset$. To \mathcal{F} we associate the set of promise problems, which are solvable with respect to \mathcal{F} , i.e. we consider $\mathbf{promise}(\mathcal{F}) = \{(A, B) \mid A \cap B = \emptyset \text{ and } \exists Q \in \mathcal{F}^{\mathbf{dc}} : A \subseteq Q \text{ and } B \subseteq Q^c\}$. We collect some elementary facts about $\mathbf{promise}(\mathcal{F})$, which follow more or less directly by the definition, especially by using the laws of De Morgan and distributivity.

Proposition 4.1.

- (1) $(A, B) \in \mathbf{promise}(\mathcal{F}) \Leftrightarrow (B, A) \in \mathbf{promise}(\mathcal{F})$.
- (2) $B' \subseteq B$ and $(A, B) \in \mathbf{promise}(\mathcal{F}) \Rightarrow (A, B') \in \mathbf{promise}(\mathcal{F})$.

- (3) $A \in \mathcal{F}^{dc}$ and $A \cap B = \emptyset \Rightarrow (A, B) \in \mathbf{promise}(\mathcal{F})$.
- (4) $A \in \mathcal{F}^{dc} \Leftrightarrow (A, A^c) \in \mathbf{promise}(\mathcal{F})$.
- (5) $\mathcal{F}' \subseteq \mathcal{F} \Rightarrow \mathbf{promise}(\mathcal{F}') \subseteq \mathbf{promise}(\mathcal{F})$.
- (6) $\mathbf{promise}(\mathcal{F}) = \mathbf{promise}(\mathcal{F}^{co}) = \mathbf{promise}(\mathcal{F}^{dc})$.
- (7) If $\mathcal{F} = (\mathcal{F}^u)^s$ then $(A, B) \in \mathbf{promise}(\mathcal{F})$ and $(A, B') \in \mathbf{promise}(\mathcal{F}) \Rightarrow (A, B \cup B') \in \mathbf{promise}(\mathcal{F})$.
- (8) If $\mathcal{F} \pm \mathcal{V} \subseteq \mathcal{F}$ then $\forall C \in \mathcal{V} : (A, B) \in \mathbf{promise}(\mathcal{F}) \Rightarrow (A \cup C, B \cap C^c) \in \mathbf{promise}(\mathcal{F})$.

Example 4.2. Consider $X = \{a, b\}$ and the languages $A = \{a^n b^n | n > 0\}$ and $B = \{a^n b^m | n, m > 0 \text{ and } n \neq m\}$. Then $A, B, A^c \in \mathcal{L}_{cf}(X)$ and $B \subseteq A^c$, hence $(A, B) \in \mathbf{promise}(\mathcal{L}_{cf}(X))$. We show, that $(A, B) \notin \mathbf{promise}(\mathcal{L}_{reg}(X))$. Suppose the contrary, i.e. a $Q \in \mathcal{L}_{reg}(X)$ exists with $A \subseteq Q$ and $B \subseteq Q^c$. Consider a word $w_0 = a^n b^n$, where n is sufficiently large. By the pumping lemma for regular sets $u, v, w \in X^*, w \neq \mathbf{1}$ exist with $w_0 = uvw$, $|uw| \leq n$ and $uw^k v \in Q$ for all $k \geq 0$. But then $uw = a^i$ for some $1 \leq i \leq n$ and therefore $uw^2 v = a^{n+|w|} b^n \in B \cap Q$. In total, $B \cap Q \neq \emptyset$ and we get a contradiction.

The following criterion asserts for a promise problem (A, B) the existence of a nontrivial solvable subproblem.

Lemma 4.3. Let $\mathcal{V} \subseteq \mathcal{F}$ and $\mathcal{F} \pm \mathcal{V} \subseteq \mathcal{F}$. Then for all $A, B \in \mathcal{F} \setminus \mathbf{fin}(S)$ with $A \notin \mathbf{cohesive}(\mathcal{V})$ a $Q \in \mathcal{V}^{dc}$ exists, such that $A \cap Q, B \cap Q^c \in \mathcal{F} \setminus \mathbf{fin}(S)$ and $(A \cap Q, B \cap Q^c) \in \mathbf{promise}(\mathcal{V})$.

Proof. Let A, B be given according to the assumption. Since $A \notin \mathbf{cohesive}(\mathcal{V})$, $Q \in \mathcal{V}^{dc}$ exists with $A \cap Q, A \cap Q^c \notin \mathbf{fin}(S)$. But then $B \cap Q$ or $B \cap Q^c$ must be infinite, because otherwise $B = (B \cap Q) \cup (B \cap Q^c) \in \mathbf{fin}(S)$. If $B \cap Q^c \notin \mathbf{fin}(S)$, $A \cap Q, B \cap Q^c \in \mathcal{F} \setminus \mathbf{fin}(S)$ and $(A \cap Q, B \cap Q^c) \in \mathbf{promise}(\mathcal{V})$. If $B \cap Q \notin \mathbf{fin}(S)$, $A \cap Q^c, B \cap Q \in \mathcal{F} \setminus \mathbf{fin}(S)$ and $(A \cap Q^c, B \cap Q) \in \mathbf{promise}(\mathcal{V})$. \square

It is interesting to look at $\mathcal{L}_{r.e.}(X)$. Consider a promise problem (A, B) with $A \in \mathcal{L}_{r.e.}(X) \setminus \mathbf{fin}(X^*)$. Then $C \subseteq A$ exists with $C \in \mathcal{L}_{rec}(X) \setminus \mathbf{fin}(X^*)$. Hence, $(C, B) \in \mathbf{promise}(\mathcal{L}_{rec}(X)) = \mathbf{promise}(\mathcal{L}_{r.e.}(X))$, since $B \subseteq A^c \subseteq C^c$. In contrast to this fact, there exists a promise problem (A, B) with $A, B \in \mathcal{L}_{r.e.}(X)$ and $(A, B) \notin \mathbf{promise}(\mathcal{L}_{r.e.}(X))$ (cf. exercise 5-34. in [8]). But if $A, B \in \mathcal{L}_{r.e.}(X)^{co}$, then $(A, B) \in \mathbf{promise}(\mathcal{L}_{r.e.}(X)^{co})$ (cf. exercise 5-33. in [8]).

We conclude this section looking at left translations.

Lemma 4.4. Let \mathcal{L} be ltr-cancellative, $\mathcal{L} = \mathcal{L}^{ltr}$ and $\mathcal{L} \pm \mathcal{L}_{reg}(X) \subseteq \mathcal{L}$. Then for all $A, B \subseteq X^*, w \in X^* : (A, B) \in \mathbf{promise}(\mathcal{L}) \Leftrightarrow (wA, wB) \in \mathbf{promise}(\mathcal{L})$.

Proof. Suppose $(A, B) \in \mathbf{promise}(\mathcal{L})$. Then $Q \in \mathcal{L}^{dc}$ exists with $A \subseteq Q$ and $B \subseteq Q^c$. But then $wA \subseteq wQ$ and $wB \subseteq wQ^c \subseteq (wQ)^c = wQ^c \cup (wX^*)^c$. Since $\mathcal{L} = \mathcal{L}^{ltr}$ and $\mathcal{L} \pm \mathcal{L}_{reg}(X) \subseteq \mathcal{L}$, we get $wQ, (wQ)^c \in \mathcal{L}$.

Conversely, suppose $(wA, wB) \in \mathbf{promise}(\mathcal{L})$. Then we find a $Q \in \mathcal{L}^{dc}$ with $wA \subseteq Q$ and $wB \subseteq Q^c$. But then $wA \subseteq Q \cap wX^* = wQ'$ and $wB \subseteq Q^c \cap wX^* = wQ''$. Since $wQ' \cup wQ'' = (Q \cap wX^*) \cup (Q^c \cap wX^*) = wX^*$ and $wQ' \cap wQ'' = \emptyset$,

$Q'' = Q'^c$. Hence, $A \subseteq Q'$ and $B \subseteq Q'^c$. Again by the closure properties of \mathcal{L} we get $Q', Q'^c \in \mathcal{L}$. \square

5. UNSOLVABILITY OF PROMISE PROBLEMS AND COHESIVENESS

The structure of promise problems is heavily influenced by cohesiveness, more precisely if cohesiveness can be connected to a promise problem (A, B) , then it is not solvable.

Theorem 5.1. *If \mathcal{F} is closed under finite variation and $A \cap B = \emptyset$, then the following statements are equivalent:*

- (1) $A, B \notin \mathbf{fin}(S)$ and $A \cup B \in \mathbf{cohesive}(\mathcal{F})$.
- (2) $A, B \in \mathbf{cohesive}(\mathcal{F})$ and $(A, B) \notin \mathbf{promise}(\mathcal{F})$.

Proof. (1) \Rightarrow (2): Let $A, B \notin \mathbf{fin}(S)$ and $A \cup B \in \mathbf{cohesive}(\mathcal{F})$, then by prop.2.7. $A, B \in \mathbf{cohesive}(\mathcal{F})$. Suppose to the contrary that $(A, B) \in \mathbf{promise}(\mathcal{F})$. Then $Q \in \mathcal{F}^{\text{dc}}$ exists with $A \subseteq Q$ and $B \subseteq Q^c$. But then $A \subseteq (A \cup B) \cap Q \notin \mathbf{fin}(S)$ and $B \subseteq (A \cup B) \cap Q^c \notin \mathbf{fin}(S)$. This contradicts $A \cup B \in \mathbf{cohesive}(\mathcal{F})$.

(2) \Rightarrow (1): Let $A, B \in \mathbf{cohesive}(\mathcal{F})$ and $(A, B) \notin \mathbf{promise}(\mathcal{F})$. Suppose that $A \cup B \notin \mathbf{cohesive}(\mathcal{F})$, i.e. a $Q \in \mathcal{F}^{\text{dc}}$ exists with $(A \cup B) \cap Q, (A \cup B) \cap Q^c \notin \mathbf{fin}(S)$. Let $A_1 = A \cap Q, B_1 = B \cap Q, A_2 = A \cap Q^c$ and $B_2 = B \cap Q^c$. Then we get the following two cases:

Case 1: $A_{1,2} \notin \mathbf{fin}(S)$ or $B_{1,2} \notin \mathbf{fin}(S)$. Then $A = A_1 \cup A_2 \notin \mathbf{cohesive}(\mathcal{F})$ or $B = B_1 \cup B_2 \notin \mathbf{cohesive}(\mathcal{F})$ - a contradiction.

Case 2: $A_1, B_2 \notin \mathbf{fin}(S)$ and $A_2, B_1 \in \mathbf{fin}(S)$ or $A_2, B_1 \notin \mathbf{fin}(S)$ and $A_1, B_2 \in \mathbf{fin}(S)$. Since $(A_1, B_2), (A_2, B_1) \in \mathbf{promise}(\mathcal{F})$, we can apply prop.4.1.(8) for $\mathcal{V} = \mathbf{fin}(S)$ and obtain $(A, B) \in \mathbf{promise}(\mathcal{F})$ - a contradiction, again. \square

We can now characterize those $A, B \in \mathbf{cohesive}(\mathcal{F})$ with $A \cup B \in \mathbf{cohesive}(\mathcal{F})$.

Theorem 5.2. *If \mathcal{F} is closed under finite variation and $A, B \in \mathbf{cohesive}(\mathcal{F})$, then the following statements are equivalent:*

- (1) $A \cup B \in \mathbf{cohesive}(\mathcal{F})$
- (2) $(A \setminus B, B) \notin \mathbf{promise}(\mathcal{F})$ or $A \cap B \notin \mathbf{fin}(S)$.

Proof. (1) \Rightarrow (2): Let $A \cup B \in \mathbf{cohesive}(\mathcal{F})$ and suppose $A \cap B \in \mathbf{fin}(S)$. Then $A \setminus B, B \notin \mathbf{fin}(S)$ and $A \setminus B, B \in \mathbf{cohesive}(\mathcal{F})$ by prop.4.1.(8). Clearly, $A \cup B = (A \setminus B) \cup B$ and $A \setminus B \cap B = \emptyset$. Hence by th.5.1 $(A \setminus B, B) \notin \mathbf{promise}(\mathcal{F})$.

(2) \Rightarrow (1): Conversely, we have to consider two cases. First suppose that $A \cap B \in \mathbf{fin}(S)$ and $(A \setminus B, B) \notin \mathbf{promise}(\mathcal{F})$. By assumption and prop.2.7. $A \setminus B, B \in \mathbf{cohesive}(\mathcal{F})$. Hence, $A \cup B = A \setminus B \cup B \in \mathbf{cohesive}(\mathcal{F})$ by th.5.1. If $A \cap B \notin \mathbf{fin}(S)$, then by le.2.8. $A \cup B \in \mathbf{cohesive}(\mathcal{F})$. \square

Th.5.1. deals essentially with "unsolvability cores" of promise problems for \mathcal{F} . This leads to the following definition:

Definition 5.3. (A, B) is a *core* of \mathcal{F} ($(A, B) \in \mathbf{core}(\mathcal{F})$) if and only if $A, B \notin \mathbf{fin}(S)$, $A \cap B = \emptyset$ and for all $A' \subseteq A, B' \subseteq B, A', B' \notin \mathbf{fin}(S) : (A', B') \notin \mathbf{promise}(\mathcal{F})$.

Proposition 5.4.

- (1) $\mathbf{core}(\mathcal{F}) = \mathbf{core}(\mathcal{F}^{co}) = \mathbf{core}(\mathcal{F}^{dc})$
- (2) $(A, B) \in \mathbf{core}(\mathcal{F}) \Leftrightarrow (B, A) \in \mathbf{core}(\mathcal{F})$.
- (3) $(A, B) \in \mathbf{core}(\mathcal{F}), A' \subseteq A, B' \subseteq B$ and $A', B' \notin \mathbf{fin}(S) \Rightarrow (A', B') \in \mathbf{core}(\mathcal{F})$.

Now, we want to show, that the condition " $A \cup B$ is \mathcal{F} -cohesive" characterizes completely the cores of \mathcal{F} . The following lemma offers a property of cores, which is similar to the definition of cohesive sets (def.2.1).

Lemma 5.5. *If $A, B \notin \mathbf{fin}(S)$ and $A \cap B = \emptyset$ then the following statements are equivalent:*

- (1) $(A, B) \in \mathbf{core}(\mathcal{F})$
- (2) $\forall Q \in \mathcal{F}^{dc} : (A \cap Q \notin \mathbf{fin}(S) \Leftrightarrow B \cap Q^c \in \mathbf{fin}(S))$

Proof. (1) \Rightarrow (2) : Let $(A, B) \in \mathbf{core}(\mathcal{F})$. Consider $Q \in \mathcal{F}^{dc}$ with $A' = A \cap Q \notin \mathbf{fin}(S)$. Suppose to the contrary that $B' = B \cap Q^c \notin \mathbf{fin}(S)$. Then $(A', B') \in \mathbf{promise}(\mathcal{F})$, since $A' \subseteq Q$ and $B' \subseteq Q^c$. This contradicts $(A, B) \in \mathbf{core}(\mathcal{F})$. Conversely, let $B \cap Q^c \in \mathbf{fin}(S)$ and suppose $A \cap Q \in \mathbf{fin}(S)$. Then $A' = A \cap Q^c$ and $B' = B \cap Q$ are infinite. Furthermore $A' \subseteq Q^c$ and $B' \subseteq Q$. Hence, $(A', B') \in \mathbf{promise}(\mathcal{F})$ and we get again a contradiction.

(2) \Rightarrow (1) : Let the equivalence be valid for any $Q \in \mathcal{F}^{dc}$. Suppose to the contrary that $(A, B) \notin \mathbf{core}(\mathcal{F})$. Then $A' \subseteq A, B' \subseteq B$ exist with $A', B' \notin \mathbf{fin}(S)$ and $(A', B') \in \mathbf{promise}(\mathcal{F})$. Hence, we can find $Q \in \mathcal{F}^{dc}$ with $A' \subseteq Q, B' \subseteq Q^c$. But then $A' \subseteq A \cap Q$ and $B' \subseteq B \cap Q^c$, i.e. $A \cap Q, B \cap Q^c \notin \mathbf{fin}(S)$ - a contradiction to the equivalence. \square

Remark: By prop.5.4.(1) le.5.5.(2) is equivalent to: $\forall Q \in \mathcal{F}^{dc} : (B \cap Q \notin \mathbf{fin}(S) \Leftrightarrow A \cap Q^c \in \mathbf{fin}(S))$.

Theorem 5.6. *If \mathcal{F} is closed under finite variation, $A \cap B = \emptyset$ and $A, B \notin \mathbf{fin}(S)$, then the following statements are equivalent:*

- (1) $(A, B) \in \mathbf{core}(\mathcal{F})$
- (2) $A \cup B \in \mathbf{cohesive}(\mathcal{F})$

Proof. (1) \Rightarrow (2) : Let $(A, B) \in \mathbf{core}(\mathcal{F})$. Consider $Q \in \mathcal{F}^{dc}$ with $(A \cup B) \cap Q \notin \mathbf{fin}(S)$. Then $A \cap Q \notin \mathbf{fin}(S)$ or $B \cap Q \notin \mathbf{fin}(S)$. If $A \cap Q$ is finite, $B \cap Q$ must be infinite. But then $A \cap Q^c \in \mathbf{fin}(S)$ by le.5.5. and therefore $A \in \mathbf{fin}(S)$, which contradicts the assumption $A \notin \mathbf{fin}(S)$. Hence, $A \cap Q$ must be infinite. By the same reason $B \cap Q$ must be infinite, too. But then $B \cap Q^c, A \cap Q^c \in \mathbf{fin}(S)$ by le.5.5. and therefore $(A \cup B) \cap Q^c \in \mathbf{fin}(S)$. In total $A \cup B \in \mathbf{cohesive}(\mathcal{F})$.

(2) \Rightarrow (1) : Suppose that $A \cup B \in \mathbf{cohesive}(\mathcal{F})$. Let $A' \subseteq A, B' \subseteq B$ and $A', B' \notin \mathbf{fin}(S)$. Then $A' \cap B' = \emptyset$ and $A' \cup B' \subseteq A \cup B$, i.e. $A' \cup B' \in \mathbf{cohesive}(\mathcal{F})$. But then $(A', B') \notin \mathbf{promise}(\mathcal{F})$ by th.5.1. and therefore $(A, B) \in \mathbf{core}(\mathcal{F})$. \square

Corollary 5.7. *If \mathcal{F} is closed under finite variation, $A, B, C \notin \mathbf{fin}(S)$, $B \subseteq C$ and $A \cap C = \emptyset$, then $(A, B) \in \mathbf{core}(\mathcal{F})$ and $C \in \mathbf{cohesive}(\mathcal{F})$ implies $(A, C) \in \mathbf{core}(\mathcal{F})$.*

Proof. Let A, B, C be given according to the assumption. Consider $(A, B) \in \mathbf{core}(\mathcal{F})$ and $C \in \mathbf{cohesive}(\mathcal{F})$. Then $A \cup B \in \mathbf{cohesive}(\mathcal{F})$ by th.5.6. Since $(A \cup B) \cap C = B \notin \mathbf{fin}(S)$, we get $(A \cup B) \cup C = A \cup C \in \mathbf{cohesive}(\mathcal{F})$ by le.2.8. In conclusion $(A, C) \in \mathbf{core}(\mathcal{F})$ by th.5.6. \square

Moreover, we obtain the transitivity of cores.

Corollary 5.8. *If \mathcal{F} is closed under finite variation and $A \cap B = A \cap C = B \cap C = \emptyset$, then $(A, B) \in \mathbf{core}(\mathcal{F})$ and $(B, C) \in \mathbf{core}(\mathcal{F})$ implies $(A, C) \in \mathbf{core}(\mathcal{F})$.*

Proof. Let A, B, C be given according to the assumption. Let $(A, B) \in \mathbf{core}(\mathcal{F})$ and $(B, C) \in \mathbf{core}(\mathcal{F})$. Then $B \cup C \in \mathbf{cohesive}(\mathcal{F})$ by th.5.6. Hence $C \in \mathbf{cohesive}(\mathcal{F})$ by prop.2.7. But now, $(A, B \cup C) \in \mathbf{core}(\mathcal{F})$ by cor.5.7. and therefore $(A, C) \in \mathbf{core}(\mathcal{F})$ by prop.5.4.(2). \square

Combining th.5.6 with le.2.3. we get

Corollary 5.9. *If \mathcal{F} is closed under finite variation and $\mathcal{F} = \mathcal{F}^{cc}$, then $\mathbf{core}(\mathcal{F}) = \mathbf{core}(\mathcal{F}^b)$.*

In contrast to the transitivity of cores, the property not belonging to $\mathbf{promise}(\mathcal{F})$ is transitive only with an additional condition.

Lemma 5.10. *Let \mathcal{F} be closed under finite variation and $A \cap B = A \cap C = B \cap C = \emptyset$. If $B \in \mathbf{cohesive}(\mathcal{F})$, then $(A, B) \notin \mathbf{promise}(\mathcal{F})$ and $(B, C) \notin \mathbf{promise}(\mathcal{F})$ implies $(A, C) \notin \mathbf{promise}(\mathcal{F})$.*

Proof. Suppose that $(A, B) \notin \mathbf{promise}(\mathcal{F})$, $(B, C) \notin \mathbf{promise}(\mathcal{F})$ and $(A, C) \in \mathbf{promise}(\mathcal{F})$. Let $Q \in \mathcal{F}^{cd}$ with $A \subseteq Q$ and $C \subseteq Q^c$. Since $B \in \mathbf{cohesive}(\mathcal{F})$, $B \cap Q$ or $B \cap Q^c$ has to be finite. By symmetry we can assume, that $D = B \cap Q \in \mathbf{fin}(S)$. But then, $A \cup D \subseteq Q$ and $C \cup (B \cap D^c) \subseteq Q^c$, i.e. $(A \cup D, C \cup (B \cap D^c)) \in \mathbf{promise}(\mathcal{F})$. Thus, $(A \cup D, B \cap D^c) \in \mathbf{promise}(\mathcal{F})$, too and we can apply prop.4.1.(8) and conclude, that $(A, B) \in \mathbf{promise}(\mathcal{F})$ - a contradiction. \square

As shown in [1] by a marking technique this kind of transitivity is not valid in the general case. The same technique is used in

Example 5.11. Let $X = \{a, b\}$. Consider a language family \mathcal{L} satisfying the condition of le.4.4. Choose A with $A, A^c \notin \mathcal{L}$. Then $(A, A^c) \notin \mathbf{promise}(\mathcal{L})$ and by le.4.4 $(xA, xA^c) \notin \mathbf{promise}(\mathcal{L})$ for any $x \in X$. Hence $(aA, aA^c \cup bA^c), (aA^c \cup bA^c, bA) \notin \mathbf{promise}(\mathcal{L})$. But $(aA, bA) \in \mathbf{promise}(\mathcal{L})$.

Next we want to show, that under some closure condition for \mathcal{F} any $(A, B) \notin \mathbf{promise}(\mathcal{F})$ contains a core of \mathcal{F} . We shall use a construction similar to the construction in the proof of the Dekker-Myhill theorem. For this purpose the following lemma is crucial.

Lemma 5.12. *If $\mathcal{F}^u = \mathcal{F} = \mathcal{F}^s$ and $(A, B) \notin \mathbf{promise}(\mathcal{F})$, then for all $Q \in \mathcal{F}^{dc}$: $(A \cap Q, B \cap Q) \notin \mathbf{promise}(\mathcal{F})$ or $(A \cap Q^c, B \cap Q^c) \notin \mathbf{promise}(\mathcal{F})$.*

Proof. Suppose to the contrary a $Q \in \mathcal{F}^{dc}$ exists, such that $(A \cap Q, B \cap Q) \in \mathbf{promise}(\mathcal{F})$ and $(A \cap Q^c, B \cap Q^c) \in \mathbf{promise}(\mathcal{F})$. Then we can find $Q_{1,2} \in \mathcal{F}^{dc}$ with $A \cap Q \subseteq Q_1, B \cap Q \subseteq Q_1^c$ and $A \cap Q^c \subseteq Q_2, B \cap Q^c \subseteq Q_2^c$. Now, $A = (A \cap Q) \cup (A \cap Q^c) \subseteq (Q_1 \cap Q) \cup (Q_2 \cap Q^c)$ and $B = (B \cap Q) \cup (B \cap Q^c) \subseteq (Q_1^c \cap Q) \cup (Q_2^c \cap Q^c)$. Let $Q_A = (Q_1 \cap Q) \cup (Q_2 \cap Q^c)$ and $Q_B = (Q_1^c \cap Q) \cup (Q_2^c \cap Q^c)$. Then $Q_A \cup Q_B = Q \cup Q^c = S$ and $Q_A \cap Q_B = \emptyset$. Hence, $Q_B = Q_A^c$. Since $A \subseteq Q_A, B \subseteq Q_B = Q_A^c$ and $Q_A, Q_B \in \mathcal{F}$ we get by the closure properties of \mathcal{F} in total $(A, B) \in \mathbf{promise}(\mathcal{F})$ - a contradiction. \square

Theorem 5.13. *If \mathcal{F} is denumerable, closed under finite variation and $\mathcal{F}^u = \mathcal{F} = \mathcal{F}^s$, then for all (A, B) with $A \cap B = \emptyset$ and $(A, B) \notin \mathbf{promise}(\mathcal{F})$, $A' \subseteq A, B' \subseteq B$ exist, such that $(A', B') \in \mathbf{core}(\mathcal{F})$.*

Proof. Let $e_{\mathcal{F}} : \mathbb{N}_0 \rightarrow 2^S$ with $e_{\mathcal{F}}(\mathbb{N}_0) = \mathcal{F}$ and $(A, B) \notin \mathbf{promise}(\mathcal{F})$. Then we construct the following sequence of pairs (A_n, B_n) for $n \geq 0$ inductively by

$$\begin{aligned} (A_0, B_0) &= (A, B) \\ (A_{n+1}, B_{n+1}) &= \mathbf{if} (A_n \cap e_{\mathcal{F}}(n), B_n \cap e_{\mathcal{F}}(n)) \notin \mathbf{promise}(\mathcal{F}) \\ &\quad \mathbf{then} (A_n \cap e_{\mathcal{F}}(n), B_n \cap e_{\mathcal{F}}(n)) \\ &\quad \mathbf{else} (A_n \cap e_{\mathcal{F}}(n)^c, B_n \cap e_{\mathcal{F}}(n)^c) \mathbf{fi}. \end{aligned}$$

Assertion 1: $\forall n \geq 0 : A_{n+1} \subseteq A_n, B_{n+1} \subseteq B_n$ and $(A_n, B_n) \notin \mathbf{promise}(\mathcal{F})$.

Clearly, $A_{n+1} \subseteq A_n$ and $B_{n+1} \subseteq B_n$ for $n \geq 0$ follows directly from the definition. The second part of the assertion is proven by induction on n . If $n = 0$ then $(A_0, B_0) = (A, B) \notin \mathbf{promise}(\mathcal{F})$ by assumption. Consider (A_{n+1}, B_{n+1}) . If $(A_n \cap e_{\mathcal{F}}(n), B_n \cap e_{\mathcal{F}}(n)) \notin \mathbf{promise}(\mathcal{F})$, nothing is to prove. Suppose $(A_n \cap e_{\mathcal{F}}(n), B_n \cap e_{\mathcal{F}}(n)) \in \mathbf{promise}(\mathcal{F})$. Since $\mathcal{F}^u = \mathcal{F} = \mathcal{F}^s$, we get by le.5.12. $(A_{n+1}, B_{n+1}) = (A_n \cap e_{\mathcal{F}}(n)^c, B_n \cap e_{\mathcal{F}}(n)^c) \notin \mathbf{promise}(\mathcal{F})$.

Assertion 2: $\forall n \geq 0 \exists k \geq n : A_k \subset A_n$ and $B_k \subset B_n$.

Assume $n \geq 0$ exists with $A_j = A_n$ for all $j \geq n$. Since $A_n \notin \mathbf{fin}(S)$, $x, y \in A_n$ exist with $x \neq y$. Since \mathcal{F} is closed under finite variation, $x \in e_{\mathcal{F}}(m)$ and $y \in e_{\mathcal{F}}(m)^c$ for some m . Furthermore $m \geq n$, otherwise x and y can not be both in A_n . Hence, by construction either $x \notin A_{m+1}$ or $y \notin A_{m+1}$, i.e. $A_{m+1} \neq A_n$, while on the other side by our assumption $A_{m+1} = A_m = A_n$ - a contradiction. Analogously, $m' \geq n$ exists with $B_{m'} \subset B_n$. Choosing $k = \mathbf{max}(m, m')$ we get the result by ass.1.

On the basis of ass.2 a function $g : \mathbb{N}_0 \rightarrow \mathbb{N}_0$ exists with $A_{g(i+1)} \subset A_{g(i)} \subset A_i$ and $B_{g(i+1)} \subset B_{g(i)} \subset B_i$ for any $i \geq 0$. But then two sequences a_i and b_i exist with $a_i \in A_{g(i)} \setminus A_{g(i+1)}$ and $b_i \in B_{g(i)} \setminus B_{g(i+1)}$ ($i \geq 0$) and the property: $0 \leq i < j \Rightarrow a_i \neq a_j$ and $b_i \neq b_j$. Hence, the two sets $A' = \{a_i | i \geq 0\}$ and $B' = \{b_i | i \geq 0\}$ are both infinite. Furthermore, $A' \cap B' = \emptyset$, since $A' \subseteq A, B' \subseteq B$ and $A \cap B = \emptyset$.

Now, we can show that $A' \cup B' \in \mathbf{cohesive}(\mathcal{F})$. Then $A' \cup B' \in \mathbf{core}(\mathcal{F})$ follows directly by th.5.6. completing the proof of the theorem.

Consider $Q \in \mathcal{F}^{\text{dc}}$, i.e. $Q = \mathbf{e}_{\mathcal{F}}(m)$ for some $m \geq 0$. Then $(A_{m+1}, B_{m+1}) = (A_m \cap \mathbf{e}_{\mathcal{F}}(m), B_m \cap \mathbf{e}_{\mathcal{F}}(m))$ or $(A_{m+1}, B_{m+1}) = (A_m \cap \mathbf{e}_{\mathcal{F}}(m)^c, B_m \cap \mathbf{e}_{\mathcal{F}}(m)^c)$. Hence, by definition of g :

- (1) $A_{g(m+1)} \cup B_{g(m+1)} \subset A_{m+1} \cup B_{m+1} \subset \mathbf{e}_{\mathcal{F}}(m)$ or
- (2) $A_{g(m+1)} \cup B_{g(m+1)} \subset A_{m+1} \cup B_{m+1} \subset \mathbf{e}_{\mathcal{F}}(m)^c$.

Consider the first case. Observe that $A' \cap A_{g(k)} = \{a_i | i \geq k\}$ and $B' \cap B_{g(k)} = \{b_i | i \geq k\}$, i.e. almost all a_i belong to $A' \cap A_{g(k)}$ and almost all b_i belong to $B' \cap B_{g(k)}$ for any $k \geq 0$. Noticing $A' \cap B = \emptyset = A \cap B'$, we get

$$\begin{aligned} (A' \cup B') \cap Q &= (A' \cup B') \cap \mathbf{e}_{\mathcal{F}}(m) \\ &= (A' \cup B') \cap (A_{g(m+1)} \cup B_{g(m+1)}) \\ &= (A' \cap A_{g(m+1)}) \cup (B' \cap B_{g(m+1)}) \\ &= (A' \cap B') \setminus C \end{aligned}$$

for some finite set C , i.e. $(A' \cup B') \cap Q^c \in \mathbf{fin}(S)$. Completely analogously, we find in the second case $(A' \cup B') \cap Q \in \mathbf{fin}(S)$. Hence, $A' \cup B' \in \mathbf{cohesive}(\mathcal{F})$. \square

Corollary 5.14. *If \mathcal{F} is denumerable, closed under finite variation and $\mathcal{F} = \mathcal{F}^{cc}$, then for all $(A, B) \notin \mathbf{promise}(\mathcal{F})$, $A' \subseteq A, B' \subseteq B$ exist with $(A', B') \in \mathbf{core}(\mathcal{F})$.*

Proof. Since $\mathbf{core}(\mathcal{F}) = \mathbf{core}(\mathcal{F}^{\text{b}})$ by cor.5.9., the conditions of th.5.13. are met for \mathcal{F}^{b} . Hence, we find $A' \subseteq A, B' \subseteq B$ with $(A', B') \in \mathbf{core}(\mathcal{F}^{\text{b}}) = \mathbf{core}(\mathcal{F})$. \square

The following example shows, that the closure conditions of th.5.13. respectively cor.5.14. are necessary. If we look for example at the family of contextfree languages which is closed neither under intersection nor under complement, we find unsolvable promise problems without cores.

Example 5.15. Consider $X = \{a, b, c\}$. For $x \in X$ and $w \in X^*$ let $|w|_x$ denote the number of occurrences of x in w . Then define for $x, y \in X$ with $x \neq y$: $L_{x,y} = \{w \in X^* | |w|_x \neq |w|_y\}$. $L_{x,y} \in \mathcal{L}_{\text{cf}}(X)$, moreover, $L_{x,y}$ is a deterministic contextfree language, hence $L_{x,y}^c \in \mathcal{L}_{\text{cf}}(X)$. Consider $A = L_{a,b} \cup L_{b,c} \cup L_{c,a}$ and $B = A^c = \{w \in X^* | |w|_a = |w|_b = |w|_c\} = L_{a,b}^c \cap L_{b,c}^c \cap L_{c,a}^c$. Then $A \in \mathcal{L}_{\text{cf}}(X), B \notin \mathcal{L}_{\text{cf}}(X), B \in \mathcal{L}_{\text{cf}}(X)^{\text{co}}$ and $B \in \mathcal{L}_{\text{cf}}(X)^{\text{s}}$. This shows $(A, B) \notin \mathbf{promise}(\mathcal{L}_{\text{cf}}(X)), (A, B) \in \mathbf{promise}(\mathcal{L}_{\text{cf}}(X)^{\text{co}})$ and $(A, B) \in \mathbf{promise}(\mathcal{L}_{\text{cf}}(X)^{\text{s}})$.

Now, suppose $A', B' \notin \mathbf{fin}(S)$ exist with $A' \subseteq A, B' \subseteq B$ and $(A', B') \in \mathbf{core}(\mathcal{L}_{\text{cf}}(X))$. Since A' is infinite, $A' \cap L_{x,y}$ is infinite for at least one of the pairs (x, y) . Assume without loss of generality $x = a$ and $y = b$. But then, $(A' \cap L_{a,b}, B') \in \mathbf{core}(\mathcal{L}_{\text{cf}}(X))$ and therefore $(A' \cap L_{a,b}, B') \notin \mathbf{promise}(\mathcal{L}_{\text{cf}}(X))$. On the other side, $A' \cap L_{a,b} \subseteq L_{a,b}$ and $B' \subseteq B \subseteq L_{a,b}^c$, i.e. $(A' \cap L_{a,b}, B') \in \mathbf{promise}(\mathcal{L}_{\text{cf}}(X))$ and we get a contradiction.

We get one further corollary of theorem 5.13. For \mathcal{F} and $A \notin \mathbf{fin}(S)$ define $\mathbf{core}(A, \mathcal{F}) = \{B | B \notin \mathbf{fin}(S), A \cap B = \emptyset \text{ and } \forall B' \subseteq B, B' \notin \mathbf{fin}(S) : (A, B') \notin \mathbf{promise}(\mathcal{F})\}$.

Corollary 5.16. *If \mathcal{F} is denumerable, closed under finite variation and $\mathcal{F}^u = \mathcal{F} = \mathcal{F}^s$, then for all $A, A^c \notin \mathcal{F} \cup \mathbf{fin}(S)^{cc}$: $\mathbf{core}(A, \mathcal{F}) \neq \emptyset$.*

Proof. By prop.4.1(4) $(A, A^c) \notin \mathbf{promise}(\mathcal{F})$. By th.5.13. $A' \subseteq A, B' \subseteq A^c$ exist with $(A', B') \in \mathbf{core}(\mathcal{F})$. Then for any $B'' \subseteq B$ with $B'' \notin \mathbf{fin}(S)$, $(A', B'') \notin \mathbf{promise}(\mathcal{F})$ and therefore $(A, B'') \notin \mathbf{promise}(\mathcal{F})$. \square

6. COMPLEXITY CORES

We have seen that under the conditions of cor.5.16. $\mathbf{core}(A, \mathcal{F}) \neq \emptyset$, provided there exists a B with $(A, B) \notin \mathbf{promise}(\mathcal{F})$. We can improve the result under the same assumption by connecting the elements of $\mathbf{core}(A, \mathcal{F})$ to the hard cores (of complexity classes) introduced in a general form by Book-Du [2]. For \mathcal{F} and A define $\mathcal{F}(A) = \{Q \in \mathcal{F} \mid Q \subseteq A\}$.

Definition 6.1. (see [2]) B is a \mathcal{F} -hardcore of A if and only if $B \notin \mathbf{fin}(S)$ and for all $C \in \mathcal{F}(A)$: $B \cap C \in \mathbf{fin}(S)$. If additionally $B \subseteq A$, then B is a proper \mathcal{F} -hardcore of A .

Lemma 6.2. *If \mathcal{F} is closed under finite variation with $\mathcal{F} = \mathcal{F}^{cc}$ and $A \cap B = \emptyset$, then*

- (1) $(A, B) \notin \mathbf{promise}(\mathcal{F}) \Leftrightarrow B^c \in \mathbf{immune}(\mathcal{F}(A^c)^{co})$.
- (2) $B \in \mathbf{core}(A, \mathcal{F}) \Leftrightarrow B$ is a proper \mathcal{F} -hardcore of A^c .

Proof. We make use of the (trivial)

Assertion: $A \subseteq Q$ and $Q \in \mathcal{F} \Leftrightarrow Q \in \mathcal{F}(A^c)^{co}$.

(1) Suppose $(A, B) \notin \mathbf{promise}(\mathcal{F})$ and $B^c \notin \mathbf{immune}(\mathcal{F}(A^c)^{co})$. Then $Q \in \mathcal{F}(A^c)^{co}$ exists with $Q \notin \mathbf{fin}(S)$ and $Q \subseteq B^c$. But then $B \subseteq Q^c, A \subseteq Q$ and $Q \in \mathcal{F}^{dc}$ - a contradiction.

Conversely, if $(A, B) \in \mathbf{promise}(\mathcal{F})$, a $Q \in \mathcal{F}^{dc}$ exists with $A \subseteq Q, B \subseteq Q^c$. But then, $Q \in \mathcal{F}(A^c)^{co}$ and $Q \subseteq B^c$. Since $A \notin \mathbf{fin}(S), Q \notin \mathbf{fin}(S)$, i.e. $B^c \notin \mathbf{immune}(\mathcal{F}(A^c)^{co})$.

(2) Consider $B \in \mathbf{core}(A, \mathcal{F})$. Then by definition $B \notin \mathbf{fin}(S)$ and $B \subseteq A^c$. Now, suppose that B is not a proper \mathcal{F} -hardcore of A^c . Then $Q \in \mathcal{F}(A^c)$ exists with $B \cap Q \notin \mathbf{fin}(S)$. Clearly, $B \cap Q \subseteq Q \subseteq A^c$. Moreover, $A \subseteq Q^c$ and $Q \in \mathcal{F} = \mathcal{F}^{dc}$. Hence $(A, B \cap Q) \in \mathbf{promise}(\mathcal{F})$, a contradiction to $B \in \mathbf{core}(A, \mathcal{F})$.

Conversely, consider a proper \mathcal{F} -hardcore B of A^c and suppose, that $B \notin \mathbf{core}(A, \mathcal{F})$, i.e. $(A, B') \in \mathbf{promise}(\mathcal{F})$ for some $B' \subseteq B$ with $B' \notin \mathbf{fin}(S)$. Then $Q \in \mathcal{F}^{dc}$ exists with $A \subseteq Q$ and $B \subseteq Q^c$. Since $B' \subseteq B \cap Q^c \notin \mathbf{fin}(S)$ and $B \cap Q^c \subseteq A^c$, B is not a proper \mathcal{F} -hardcore of A^c and we get a contradiction. \square

In [2] R. V. Book and D. Z. Du characterize hard cores in the following way:

Theorem 6.3. (*[2]*) *Let \mathcal{F} be a denumerable set family and $A \notin \mathbf{fin}(S)$. Then a proper \mathcal{F} -hardcore B of A exists if and only if $A \notin \mathcal{F}^u \oplus \mathbf{fin}(S)$.*

Furthermore they achieved the existence of recursive hard cores under the following conditions:

Theorem 6.4. ([2]) *If \mathcal{L} is WP-recursive, closed under finite variation and $\mathcal{L} = \mathcal{L}^u$, then for any $A \in \mathcal{L}_{\text{rec}}(X) \setminus \mathcal{L}$, a proper \mathcal{L} -hardcore $B \in \mathcal{L}_{\text{rec}}(X)$ of A exists.*

Combining le.6.2. with th.6.4. we obtain

Theorem 6.5. *If \mathcal{L} is WP-recursive and $\mathcal{L} = \mathcal{L}^b$, then for any $A \in \mathcal{L}_{\text{rec}}(X) \setminus \mathcal{L} : \text{core}(A, \mathcal{F}) \cap \mathcal{L}_{\text{rec}}(X) \neq \emptyset$.*

7. CONCLUDING REMARKS

It is natural, to consider n -dimensional promise problems (A_1, \dots, A_n) with $A_i \cap A_j = \emptyset$ and $A_i \subseteq S$ for $1 \leq i \neq j \leq n$. For a set family \mathcal{F} the promise problem is solvable if a partition (Q_1, \dots, Q_n) of S exists with $A_i \subseteq Q_i, Q_i \in \mathcal{F} (1 \leq i \leq n)$. For $n > 2$ cores of unsolvability can be characterized by cohesiveness of $A_1 \cup \dots \cup A_n$, too. But for $n = 3$ unsolvable promise problems exist, which have no cores of unsolvability ([1]).

Instead of $\mathcal{L}_{\text{reg}}(X)$ we can use a much smaller language family in the variation condition " $\mathcal{L} \pm \mathcal{L}_{\text{reg}}(X) \subseteq \mathcal{L}$ ". Let $\mathcal{L}_{\text{ltr}}(X) = \{w_1 L_1 \cup \dots \cup w_k L_k \mid k > 0, w_i \in X^*, L_i \in \mathbf{fn}(X^*)^{\text{cc}} 1 \leq i \leq k\} = ((\mathbf{fn}(X^*)^{\text{cc}})_{\text{ltr}})^u$. A lengthy and involved, but elementary proof shows, that $\mathcal{L}_{\text{ltr}}(X)$ is a boolean algebra and ltr-cancellative. Then we can use always the variation condition " $\mathcal{L} \pm \mathcal{L}_{\text{ltr}}(X) \subseteq \mathcal{L}$ ". Moreover, a stronger result than le.3.2 and its corollary is possible, namely a complete characterization of *cohesive*($\mathcal{L}_{\text{ltr}}(X)$) by sequential functions.

REFERENCES

- [1] KLAUS AMBOS-SPIES, ULRIKE BRANDT, MARTIN ZIEGLER: "Real Benefit of Promises and Advice", accepted for presentation at CiE 2013 and selected for publication in the springer LNCS volume
- [2] R.V.BOOK, DING-ZHU DU: "The Existence and Density of Generalized Complexity Cores", pp.718–730 in *JACM* vol.**34:3** (July 1987).
- [2] J.L.BALCAZAR, J.DIAZ, J.GABARRO: "*Structural Complexity I, EATCS Monographs on Theoretical Computer Science*", Springer Verlag, Heidelberg (1988).
- [4] S.EVEN, A.L.SELMAN AND Y.YACOBI: "The Complexity of Promise Problems with Applications to Public-Key Cryptography", pp.159–173 in *Information and Control* vol.**61** (1984).
- [5] ODED GOLDREICH: "On Promise-Problems", (*A Survey in Memory of Shimon Even*), Dep. of Comp.Science, Weizmann Institute of Science (2005).
- [6] MICHAEL A. HARRISON: "*Introduction to Formal Language Theory*", Addison-Wesley Publishing Company (1978).
- [7] G.ROZENBERG - A.SALOMAA EDS.: "Word, Language, Grammar", in *Handbook of Formal Languages* vol.**1**, Springer-Verlag (1997).
- [8] HARTLEY ROGERS JUN.: "*Theory of Recursive Functions and Effective Computability*", MacGraw-Hill Book Company (1967).
- [9] ROBERT I. SOARE: "*Recursively enumerable Sets and Degrees*", Springer Verlag (1987).

